

Quantum decoherence in a supersymmetric mechanical system

T. Shi¹, Z. Song¹ and C. P. Sun^{1,2,a,b}

¹ *Department of Physics, Nankai University, Tianjin 300071, China*

² *Institute of Theoretical Physics, The Chinese Academy of Sciences, Beijing, 100080, China*

Abstract

We make a novel observation about the decoherence phenomenon of the fermion in the Witten's supersymmetric (SUSY) quantum mechanical model. It is shown that, when the bosonic partner in the SUNY model is unobservable in a certain energy scale, the quantum coherence meant by the superposition of fermion states can not be preserved for a long time due to the quantum decoherence induced by the overlooked boson. This refers to a supercharge superselection rule similar to the charge superselection. We numerically calculate the decoherence factor characterizing the extent of decoherence. The obtained result shows the periodic decoherence with finite quantum jump. Some analytic results under the harmonic approximation for the superpotential can be obtained to confirm the numerical calculations.

Pacs numbers: 11.30.Pb, 03.65.Bz, 12.60.Jv

The original idea of supersymmetry (SUSY) is rooted in the generalization of the Poincaré symmetry for the relativistic quantum field theory[1]. Though the SUSY can not finally be confirmed as the successful way to unify the nature laws of particle physics, the SUSY has become a major method in non-relativistic quantum physics after its one dimensional model was built by E. Witten[2]. Its successful applications could be found in many fields in physics, such as, statistical physics and atomic physics[3]. However, the existence of the SUSY partners of boson or fermion predicted by the SUSY theory in high energy physics has not been discovered experimentally so far. The popular explanation for this fact is to emphasize that the energy scale at which SUSY particles may occur should be much higher than that in all current experiments[4]. In this paper we will take into account the fact of "no-finding" of SUSY partner from a "new angle". Our central idea comes from the quantum decoherence theory

Let us recall the superselection rule that can explain why there does not exist the long life superposition of the states of neutron and proton though they belong to a double sector of isospin. The basic understanding for this phenomenon is the isospin symmetry breaking due to the electromagnetic interaction. From the point view of quantum decoherence [5], the different ways of interaction between the electromagnetic field and the neutron and proton will force the electromagnetic field to evolve into two distinguishing states respectively and then the states of two different isospin states, the neutron and proton, entangle with them. This entanglement will result in the adiabatic decoherence [7] in a formal superposition of the state of neutron and proton. We notice that L. H. Ford has studied similar problem for the electron coherence with a detailed calculation[8]. As he showed, due to the coupling of the electron to the quantized electromagnetic field, the influences of both of photon emission and of the electromagnetic vacuum fluctuations can lead a decrease in the amplitude of the interference oscillations, a kind of quantum decoherence.

Actually, in any non-trivial SUSY theory, there must exist the coupling between the boson (fermion) and its fermion (boson) partner. If the SUSY partner of a particle is invisible in an experiment, its quantum information can be lost into the subspace for the SUSY partner. In other words, if there exists a SUSY system in nature, the observed boson (or fermion) could not be preserved in a pure state (a quantum superposition) for a long time since its unobservable SUSY partner can induce its quantum decoherence. This argument shows that the quantum decoherence can result in a superselection rule, which forbid some

state superpositions of the SUSY particles.

Now we demonstrate the above arguments by making use the simplest, but non trivial SUSY quantum mechanical model - the $N = 2$ SUSY quantum system introduced by E. Witten[2]. The Witten's SUSY model for $N = 2$ is defined with two super-charges

$$\begin{aligned} Q_1 &= \frac{1}{\sqrt{2}} \left(\frac{P}{\sqrt{2m}} \sigma_1 + W(x) \sigma_2 \right) \\ Q_2 &= \frac{1}{\sqrt{2}} \left(\frac{P}{\sqrt{2m}} \sigma_2 + W(x) \sigma_1 \right) \end{aligned} \quad (1)$$

on the Hilbert space $H = L^2(R) \otimes C^2$. Here $L^2(R)$ is the space of real-valued function and C^2 is the spin space; $\{\sigma_j \mid j = 1, 2, 3\}$ are the Pauli matrices. The real function $W(x)$ of the Cartesian coordinate x is customarily called SUSY potential. With these notations, the SUSY Hamiltonian

$$H = 2Q_1^2 = 2Q_2^2 \quad (2)$$

can be explicitly written as

$$H = \frac{P^2}{2m} + W^2(x) + \frac{\hbar}{\sqrt{2m}} W'(x) \sigma_3. \quad (3)$$

This Hamiltonian can also be expressed in the block-diagonal form in the σ_3 -picture

$$H = \begin{pmatrix} H_+ & 0 \\ 0 & H_- \end{pmatrix}. \quad (4)$$

Here, the Hamiltonian in the diagonal blocks are

$$H_{\pm} = \frac{P^2}{2m} + W^2(x) \pm \frac{\hbar}{\sqrt{2m}} W'(x) \quad (5)$$

which acting on the spatial space $L^2(R)$. In fact, in this model, we can understand the above Witten model as a composite system with coupling boson and fermion. Let us define the boson creation operated

$$b^\dagger = \sqrt{\frac{m\omega}{2\hbar}} \left(x - \frac{i}{m\omega} p \right) \quad (6)$$

and the fermion creation operator

$$f^\dagger = \sigma_+ = \frac{1}{2} (\sigma_1 + i\sigma_2) \quad (7)$$

where ω is a constant determined by the function $W(x)$. Then the above Hamiltonian can be written in the form

$$H = \hbar\omega \left(b^\dagger b + \frac{1}{2} \right) + W^2 \left(\sqrt{\frac{\hbar}{2m\omega}} (b^\dagger + b) \right) - \frac{\hbar\omega}{4} (b^\dagger + b)^2 + \frac{\hbar}{\sqrt{2m}} W' \left(\sqrt{\frac{\hbar}{2m\omega}} (b^\dagger + b) \right) (2f^\dagger f - 1). \quad (8)$$

Based on the quantum decoherence theory, we can make a crucial observation about the SUSY Hamiltonian. The SUSY Hamiltonian describes a quantum-measurement-like process[9], since the interaction part

$$H_I = \frac{\hbar}{\sqrt{2m}} W'(x) \sigma_3 \quad (9)$$

commutes with σ_3 , but does not commutes with the spatial part

$$H_0 = \frac{P^2}{2m} + W^2(x) \quad (10)$$

Thus the SUSY model a Stern-Gerlach type quantum measurement about the fermion variable. In other word, if the system is initially prepared in the different σ_3 -eigen states

$$|+\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad |-\rangle = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \quad (11)$$

the spin part acts on the spatial part with with different "forces" depicted by the different effective superpotentials[7]

$$V_\pm(x) = W^2(x) \pm \frac{\hbar}{\sqrt{2m}} W'(x). \quad (12)$$

These actions will create the quantum entanglement between the spin part and the spatial part. Actually, the SUSY coupling H_I dose not change the internal energy of the fermion part due to $[H_I, \sigma_3] = 0$, but the non-zero commutators $[H_I, H_0] \neq 0$ implies that the spatial part can record the information of fermion with the differnt forces

$$F_\pm = -2W(x)W'(x) \mp \frac{\hbar}{\sqrt{2m}} W''(x) \quad (13)$$

This is just all story from the so-called quantum nondemolition measurement[9].

To demonstrate the creation of quantum entanglement, we assume the SUSY system is initially prepared in a factorized state

$$|\psi(0)\rangle = (C_+ |+\rangle + C_- |-\rangle) \otimes |\phi\rangle \quad (14)$$

where $|\phi\rangle$ describes the initial spatial wave packet $\phi(x) = \langle x|\phi\rangle$. Driven by the SUSY Hamiltonian, this system will evolve into an entangled state

$$|\psi(t)\rangle = C_+ |+\rangle \otimes |\phi_+(t)\rangle + C_- |-\rangle \otimes |\phi_-(t)\rangle \quad (15)$$

where

$$|\phi_{\pm}(t)\rangle = \exp(-iH_{\pm}t) |\phi_{\pm}(0)\rangle \quad (16)$$

depicts the two-channel evolution of the SUSY system. However, $|\psi(t)\rangle$, in usual, is not an orthogonal decomposition (or a Schmidt decomposition) since $|\phi_+(t)\rangle$ and $|\phi_-(t)\rangle$ are not orthogonal with each other[10]. In the context of quantum decoherence theory[9], we introduced the decoherence factor

$$D(t) = \langle \phi_+(t) | \phi_-(t) \rangle \quad (17)$$

whose norm $|D(t)|$ measures the extent of orthogonalization or the entanglement. It was even pointed out that $D(t)$ also characterize the extent of decoherence of the spin part when the spatial part of SUSY is overlooked in certain situation.

Actually, if we discard the spatial variable of the SUSY system for some physical consideration, we only need the reduced density matrix

$$\rho = Tr_x(|\psi(t)\rangle \langle \psi(t)|) = |C_+|^2 |+\rangle \langle +| + |C_-|^2 |-\rangle \langle -| + (C_+ C_-^* |+\rangle \langle -| D(t) + h.c) \quad (18)$$

to deal with the physical processes concerning the spin variable only. Obviously, when $D(t) = 0$, the spin part of the SUSY system lose its all quantum coherence into the spatial part and then the reduced density matrix becomes of diagonal type, i.e.,

$$\rho \rightarrow \rho_{dig} = |C_+|^2 |+\rangle \langle +| + |C_-|^2 |-\rangle \langle -|. \quad (19)$$

It is very similar to the description of double-slit interference with "which-way" detection. If one can precisely determine which slit the particle will go through, the interference pattern disappears.

From the above arguments it can be concluded that, if we consider the spin 1/2 variable of the SUSY system as the fermionic degree of freedom, while the spatial parts as the bosonic degree of freedom, the quantum decoherence of the fermion part can be induced by overlooking the motion of the bosonic part. This implies that if we can not measure both the bosonic degrees of freedom in an experiment, the observed part (fermion) can not preserve its quantum coherence since the bosonic part act as an environment to monitor the Fermion system.

To quantitatively analyze the extent of decoherence of fermion in the SUSY system, we go the details in the calculation of the decoherence factor. Let us first to test the calculation in analytical way with the harmonic approximation. The superpotential $V_{\pm}(x)$ can be expanded in the terms of $x - x_0$ up to second order around the two equilibrium points $x_0^{\pm}$ satisfying $V'_{\pm}(x_0^{\pm}) = 0$ (Fig. 1). The result is just the harmonic approximation

$$V_{\pm}(x) = V_{\pm}(x_0^{\pm}) + \frac{1}{2}m(\omega_0^{\pm})^2(x - x_0^{\pm})^2 \quad (20)$$

where

$$m(\omega_0^{\pm})^2 = 2W'^2(x_0^{\pm}) \mp 4\frac{\sqrt{2m}}{\hbar}W^2(x_0^{\pm})W'(x_0^{\pm}) \pm \frac{\hbar}{\sqrt{2m}}W'''(x_0^{\pm}). \quad (21)$$

Furthermore, we can write down the Hamiltonian for the two diagonal blocks in terms of the creation and annihilation operators as

$$H_{\pm} = \hbar\omega_0^{\pm}b_{\pm}^{\dagger}b_{\pm} + f_{\pm}(b_{\pm}^{\dagger} + b_{\pm}) + E_0^{\pm} \quad (22)$$

where

$$\begin{aligned} f_{\pm} &= -m(\omega_0^{\pm})^2x_0^{\pm}\sqrt{\frac{\hbar}{2m\omega_0^{\pm}}} \\ E_0^{\pm} &= V_{\pm}(x_0^{\pm}) + \frac{1}{2}m(\omega_0^{\pm})^2x_0^{\pm 2}. \end{aligned} \quad (23)$$

Obviously, the above Hamiltonian depict two forced harmonic oscillators with different external forces

$$F = m(\omega_0^{\pm})^2x_0^{\pm} \quad (24)$$

They will drive the two blocks evolve along two difference trajectories[7]. The decoherence factor depends on the overlap of the two wave packets along with two trajectories. The numerical calculation of the overlapping integral shows the phenomenon of quantum jump in the progressive decoherence. The quantum coherence will revive in periodic time domain.




FIG. 1: The harmonic approximation is taken in the SUSY system. The superpotential $V_{\pm}(x)$ is expanded in the terms of $x - x_0$ up to second order around the two equilibrium points $x_0^{\pm}$.

This is substantially due to the reversibility of the Schrodinger equation for the quantum system with finite degrees of freedom. This progressive decoherence phenomenon was found even theoretically in reference [9] in 1993, and the possibility of implementing its observation in cavity QED experiment was also pointed out by us in ref.[11]. In 1997 it was also independently discussed [12] with another cavity QED setup.

Under the harmonic approximation we can obtain the results analytically. We explicitly solve the Schrodinger equations of the evolution operators $U_{\pm}(t)$ in the interaction picture

with the Wei-Norman Ansatz

$$U_{\pm}(t) = e^{ip_{\pm}(t)} \exp e^{\alpha_{\pm}(t)b_{\pm}^{\dagger}} e^{\beta_{\pm}(t)b_{\pm}} = e^{iQ_{\pm}(t)} D[A_{\pm}(t)]. \quad (25)$$

where

$$D[A_{\pm}] = \exp \left[A_{\pm}^{\dagger} b_{\pm} - A_{\pm} b_{\pm}^{\dagger} \right] \quad (26)$$

are the displacement operators generating the coherent states $|\alpha_{\pm}(t)\rangle = D[A_{\pm}(t)]|0\rangle$ and

$$\begin{aligned} iQ_{\pm}(t) &= ip_{\pm} - \frac{1}{2} |A_{\pm}(t)|^2, \\ A_{\pm}(t) &= -\alpha_{\pm}(t) = \frac{g_{\pm}}{\omega_0^{\pm}} [\exp(i\omega_0^{\pm}t) - 1]. \end{aligned} \quad (27)$$

The time-dependent coefficients $p_{\pm}(t)$, $\alpha_{\pm}(t)$ and $\beta_{\pm}(t)$ can be determined by solving the differential equations about them exactly

$$\begin{aligned} ip_{\pm} &= i \left(\frac{g_{\pm}^2}{\omega_0^{\pm}} - \frac{E_0^{\pm}}{\hbar} \right) t + \frac{g_{\pm}^2}{\omega_0^{\pm 2}} [\exp(-i\omega_0^{\pm}t) - 1] \\ \alpha_{\pm}(t) &= -\frac{g_{\pm}}{\omega_0^{\pm}} [\exp(i\omega_0^{\pm}t) - 1], \\ \beta_{\pm}(t) &= \frac{g_{\pm}}{\omega_0^{\pm}} [\exp(-i\omega_0^{\pm}t) - 1], \end{aligned} \quad (28)$$

where $g_{\pm} = f_{\pm}/\hbar$. In the position reprsetation they just are the two wave packets

$$\Psi_{\pm}(t, x) = \langle x | \alpha_{\pm}(t) \rangle \quad (29)$$

which is illustrated in Fig2.

By a straightforward calculation we can analytically obtain the decoherence factor as

$$\begin{aligned} D(t) &= \langle \Psi_+(t) | \Psi_-(t) \rangle = \langle 0 | U_+^{\dagger}(t) U_-(t) | 0 \rangle = e^{iQ_-(t) - iQ_+(t)} \times \\ &\exp \left[-\frac{1}{2} \left(A_+ W - A_+^{\dagger} Y \right) \left(A_+^{\dagger} X - A_+ V \right) - \left(A_+ W - A_+^{\dagger} Y \right) A_- - \frac{1}{2} |A_-(t)|^2 \right] \end{aligned} \quad (30)$$

where

$$\begin{aligned} V &= \frac{1}{2} \sqrt{\frac{\omega_0^+}{\omega_0^-}} + \frac{im}{2} \sqrt{\omega_0^+ \omega_0^-}, \\ W &= \frac{1}{2} \sqrt{\frac{\omega_0^+}{\omega_0^-}} - \frac{im}{2} \sqrt{\omega_0^+ \omega_0^-}, \\ X &= \frac{-i}{2m} \sqrt{\frac{1}{\omega_0^+ \omega_0^-}} - \frac{1}{2} \sqrt{\frac{\omega_0^-}{\omega_0^+}}, \\ Y &= \frac{-i}{2m} \sqrt{\frac{1}{\omega_0^+ \omega_0^-}} + \frac{1}{2} \sqrt{\frac{\omega_0^-}{\omega_0^+}}. \end{aligned} \quad (31)$$




FIG. 2: The schematic illustration of the wave packet evolves in the SUSY potential.

To illustrate the above formulism visually, a simple example

$$W = \frac{1}{\sqrt{2}}Cx^2 \quad (32)$$

is chosen here. In the following we take $m = \hbar = 1$ for simplicity. It is easy to find that

$$x_0^\pm = \pm \sqrt[3]{\frac{1}{2C}}, \omega_0^\pm = \omega_0 = 6C^2 \sqrt[3]{\frac{1}{4C^2}} \quad (33)$$

and

$$D(t) = \exp\left(-\frac{8g_\pm^2}{\omega_0^2} \sin^2 \frac{\omega_0 t}{2}\right) = \exp\left(-4\omega_0 x_0^2 \sin^2 \frac{\omega_0 t}{2}\right). \quad (34)$$




FIG. 3: The decoherence factors as function of time for $c = 0.1, 0.5, \text{ and } 1.0$.

This calculations under the harmonic approximation confirm the numerical prediction about the quantum jump in the progressive decoherence, which is plotted in Fig. 3 for $c = 0.1, 0.5, 1.0$.

Finally we conclude this paper with two remarks. 1. For the quantum decoherence phenomenon in high energy regime, we even studied the quantum decoherence in neutrino flavor oscillation caused by an environment surrounding neutrinos[13]. In this way, the Ellis, Hagelin, Nanopoulos and Srednicki (EHNS) mechanism[14] for solving the solar neutrino problem can be comprehended in a framework of the ordinary quantum mechanics. Because the two kinds of neutrinos have the different interaction the weak- coupling environment,

a microscopic model was proposed to describe the transition of two neutrino system from a pure state to a mixed state. It gives the modified formula of survival probability of neutrino oscillation with two additional time-dependent parameters. For specified environments, this result shows that the oscillating phenomena of neutrino still exist even without a mass difference in free neutrino. 2. As has been mentioned in the beginning of this paper, the SUSY was originally invoked to particle physics in search of the gainful unification of nature laws with space-time and internal symmetries. The self-consistent physics in usual works well in the four dimensional quantum field theory. Our arguments about the decoherence induced by the invisible SUSY partners seem to be limited to the non-relativistic regime and thus are not able to resolve the final problem in SUSY theory. However, we believe that our central idea can be generalized for the relativistic quantum field theories. In this sense, the facts revealed in our studies in the non-relativistic can be universal. We have to point out the possible problems in our studies that the quantitative relations between the extent of decoherence and the SUSY coupling predict the fundamental SUSY gauge theories.

We acknowledge the support of the CNSF (grant No. 90203018), the Knowledge Innovation Program (KIP) of the Chinese Academy of Sciences and the National Fundamental Research Program of China (No. 001GB309310).

[a] Electronic address: suncp@itp.ac.cn

[b] Internet www site: <http://www.itp.ac.cn/~suncp>

[1] J. Wess and B. Zumino, Nucl. Phys. **B70**, 39(1974).

[2] E.Witten, Nucl. Phys. **B188**, 513(1981); **B202**(1982), 253.

[3] G.Junker, *Supersymmetric Methods in Quantum and Statistical Physics*, (Text and Monographs in Physics, Springer-Verlag, Berlin, 1996) .

[4] Y. Shadmi and Yu. Shirman, Rev. Mod. Phys. **72**, 25-64 (2000), refs therein.

[5] E. Joos, H.D. Zeh, C. Kiefer, D. Giulini, J. Kupsch and I.O. Stamatescu, (eds.) *Decoherence and the Appearance of a Classical World in Quantum Theory*, (Berlin:Springer-Verlag 2003)

[6] W. Zurek, Rev. Mod. Phys. **75**, 715 (2003).

[7] C.P. Sun, X. F. Liu, D. L. Zhou, S. X. Yu, Phys. Rev. **A** 63,01211, (2001); Eur. Phys. J. **D** 13, 145 (2001).

- [8] L. H. Ford, Phys. Rev. **D** 47, 5571-5580 (1993)
- [9] C.P. Sun, Phys. Rev. **A** 48, 878, (1993); Chin. J. Phys. 4, 7(1994); in Quantum-Classical Correspondence, edited by D.H. Feng and B.L. Hu International Press, Somerville, MA, (1997), pp. 99–106; in Quantum Coherence and Decoherence, edited by K. Fujikawa and Y.A. Ono Elsevier Science, Amsterdam, (1996), pp. 331–334; C.P. Sun et al. Fortschr. Phys. 43, 585 (1995).
- [10] D. Bouwmeeste, A. Ekert, and A. Zeilinger (Ed.), *The Physics of Quantum Information* (Springer, Berlin, 2000).
- [11] C. P. Sun, *et.al.*, Quantum Semiclassic Opt **9**, 119 (1997).
- [12] M. Brune,*et.al.*, Phys. Rev. Lett. 77, 4887 (1996); J. M. Raimond, M. Brune and S. Haroche, Phys. Rev. Lett 79, 1964 (1997).
- [13] C. P. Sun, D.L.Zhou, Quantum decoherence effect and neutrino oscillation, hep-ph/9808334,1998
- [14] J. Ellis, J. S. Hagelin, D. V. Nanopoulos and M. Srednicki, Nucl. Phys. B 241, 381(1984).